

The role of binomial type sequences in determination identities for Bell polynomials

Miloud Mihoubi

U.S.T.H.B., Faculty of Mathematics,
B.P. 32, El-Alia, 16111, Bab Ezzouar,
Algiers, Algeria.

June 20, 2008

Abstract. Our paper deals about identities involving Bell polynomials. Some identities on Bell polynomials derived using generating function and successive derivatives of binomial type sequences. We give some relations between Bell polynomials and binomial type sequences in first part, and, we generalize the results obtained in [4] in second part.

Keywords. Partial and complete Bell polynomials. Binomial type sequences. Stirling numbers. Appell polynomials. Generating functions.

1 Introduction

Recall that the (exponential) partial Bell polynomials $B_{n,k}(x_1, x_2, \dots)$ are defined by their generating function:

$$\sum_{n=k}^{\infty} B_{n,k}(x_1, x_2, \dots) \frac{t^n}{n!} = \frac{1}{k!} \left(\sum_{m=1}^{\infty} x_m \frac{t^m}{m!} \right)^k. \quad (1)$$

and the (exponential) complete Bell polynomials $A_n(x_1, x_2, \dots)$ are given by

$$A_n(x_1, x_2, \dots) := \sum_{k=1}^n B_{n,k}(x_1, x_2, \dots) \quad \text{with} \quad A_0(x_1, x_2, \dots) := 1.$$

Comtet [3] studies the Bell polynomials and gives some basic properties for them. Some applications of Bell polynomials are given by Riordan [5] in combinatorial analysis and by S. Roman [6] in umbral calculus. Recently, by using the Lagrange inversion formula (LIF), Abbas and Bouroubi [1] give some identities for the partial Bell polynomials, and, Mihoubi [4] also gives some extensions involving to the partial and complete Bell polynomials. For any sequence $(x_n; n \geq 1)$ with $x_1 = 1$ and any natural numbers

r, s , recall that Proposition 4 in [4] gives:

$$\begin{aligned} B_{n,k} \left(B(s, s), \dots, \frac{ms}{r(m-1)+s} \frac{B((r+1)(m-1)+s, r(m-1)+s)}{\binom{(r+1)(m-1)+s}{r(m-1)+s}}, \dots \right) \\ = \binom{n}{k} \frac{sk}{r(n-k)+sk} \frac{B((r+1)(n-k)+sk, r(n-k)+sk)}{\binom{(r+1)(n-k)+sk}{r(n-k)+sk}}, \end{aligned} \quad (2)$$

where $B(n, k) := B_{n,k}(x_1, x_2, x_3, \dots)$.

Then if we put

$$Y(n, k) := \binom{n}{k} \frac{sk}{r(n-k)+sk} \frac{B((r+1)(n-k)+sk, r(n-k)+sk)}{\binom{(r+1)(n-k)+sk}{r(n-k)+sk}}, \quad (3)$$

we conclude, from (2), that the sequence $(Y(n, k))$ satisfies the equation:

$$B_{n,k}(Y(1, 1), Y(2, 1), Y(3, 1), \dots) = Y(n, k). \quad (4)$$

In general, from the definition (1), if we put $\psi(t) = \sum_{m=1}^{\infty} x_m \frac{t^m}{m!}$ and $Y(n, k) = \frac{1}{k!} D_{t=0}^n (\psi(t))^k$, then for all integers n, k , with $n \geq k \geq 1$, the sequence $(Y(n, k))$ satisfies (4). Hence, to find identities for the partial Bell polynomials, it suffices to find sequences $(Y(n, k))$ satisfy the equation (4).

Similarly to the partial Bell polynomials, for any sequence $(x_n; n \geq 1)$ with $x_1 = 1$ and any natural numbers r, s ($r \geq 1$), another relation for the complete Bell polynomials is given by Proposition 8 in [4] by:

$$A_n \left(\frac{sB(r+1, r)}{r(r+1)}, \dots, \frac{s}{nr} \frac{B((r+1)n, nr)}{\binom{(r+1)n}{nr}} \right) = \frac{s}{nr+s} \frac{B((r+1)n+s, nr+s)}{\binom{(r+1)n+s}{nr+s}} \quad (5)$$

where $B(n, k) : = B_{n,k}(x_1, x_2, x_3, \dots)$,

and if we put

$$Z(n, s) := \frac{1}{nr+s} \frac{B((r+1)n+s, nr+s)}{\binom{(r+1)n+s}{nr+s}}, \quad (6)$$

we conclude, from (5), that the sequence $(Z(n, s))$ satisfies the equation:

$$A_n(sZ(1, 0), sZ(2, 0), \dots, sZ(n, 0)) = sZ(n, s). \quad (7)$$

Hence, to find identities for the complete Bell polynomials, it suffices to find sequences $(Z(n, s))$ satisfy the equation (7). Therefore, to determine solutions for (4) and (7), we exploit the strong connection between Bell polynomials and binomial type sequences. For any binomial type sequence $(f_n(x))$, with $f_0(x) := 1$, one of such connections is given in [6, p. 82] by

$$f_n(x) = \sum_{k=1}^n B_{n,k}(x_1, x_2, \dots) x^k \quad \text{with} \quad x_n = \frac{d}{dx} f_n(x)|_{x=0}. \quad (8)$$

On the basis of the results obtained in [4] and the relation (8), we derive in this paper some interesting identities and relations related Bell polynomials and binomial type sequences.

For the next of this paper, we will denote by $D_t f(t)$, $D_t^j f(t)$, $D_{t=x} f(t)$ or $Df(x)$ and $D_{t=x}^j f(t)$, respectively, for the derivative of f , the j -th derivative of f , the derivative of f evaluated at $t = x$, and the j -th derivative of f evaluated at $t = x$.

2 Main results

For the next of this work, for any sequence $(f_n(x))$ of binomial type with $f_0(x) = 1$ and for a given real number a , we define

$$f_n(x; a) := \frac{x}{an+x} f_n(an+x) \quad \text{with} \quad f_0(x; a) = 1. \quad (9)$$

The sequence $(f_n(x; a))$ is of binomial type, see [4].

To simplify any expression below we put

$$T = T(n, k) := r(n-k) + sk, \quad R = R(n, s) := nr + s$$

and $(f_n(x))$ denotes a sequence of binomial type with $f_0(x) = 1$.

The two following theorems give some interesting relations between Bell polynomials and binomial type sequences. These relations are used to deduce several identities for partial and complete Bell polynomials as it is illustrated below. To prove these theorems, we use the following Lemma:

Lemma 1 *Let n, k be integers with $n \geq k \geq 1$ and a, α be a real numbers. We have*

$$\begin{aligned} B_{n,k}(\alpha, 2D_{z=0}(e^{\alpha z} f_1(x+z; a)), \dots, mD_{z=0}(e^{\alpha z} f_{m-1}(kx+z; a)), \dots) \\ = \binom{n}{k} D_{z=0}^k(e^{\alpha z} f_{n-k}(kx+z; a)). \end{aligned} \quad (10)$$

This identity can be replaced when $\alpha = 0$ by

$$B_{n,k}(D_x f_1(x; a), \dots, D_x f_m(x; a), \dots) = \frac{1}{k!} D_{z=0}^k f_n(kx+z; a). \quad (11)$$

Theorem 2 *Let a, x, α be real numbers and n, k, r, s be natural numbers with $n \geq k \geq 1$ and $r + s \geq 1$. Then the sequence*

$$Y(n, k) := \binom{n}{k} \frac{sk}{T} D_{z=0}^T(e^{\alpha z} f_{n-k}(Tx+z; a)) \quad (12)$$

satisfies (4). For $\alpha = 0$ the sequence $(Y(n, k))$ can be replaced by

$$Y(n, k) := \frac{n!}{k!(T+n-k)!} \frac{sk}{T} D_{z=0}^T f_{T+n-k}(Tx+z; a). \quad (13)$$

For $r = s = 0$, we put $Y(n, k) := \binom{n}{k} f_{n-k}(x; a)$.

Theorem 3 *Let a, x, α be real numbers and n, r, s be natural numbers with $n \geq 1, r \geq 1$. Then for $\alpha \neq 0$ the sequence*

$$Z(n, s) := \frac{1}{\alpha^s} \frac{1}{R} D_{z=0}^R(e^{\alpha z} f_n(Rx+z; a)) \quad (14)$$

satisfies (7), and (case $\alpha = 0$) the sequence

$$Z(n, s) := \frac{1}{(Df_1(0))^s} \frac{1}{R} \frac{n!}{(R+n)!} D_{z=0}^R f_{R+n}(Rx+z; a) \quad (15)$$

satisfies (7).

More generally, Theorem 2 can be generalized as follows:

Theorem 4 Let $(a_n; n \geq 1)$ be a real sequence; n, k, r, s, u, v be natural numbers with $n \geq k \geq 1, r+s \geq 1$ and $x, a, \alpha, \beta, \lambda$ be real numbers. Then the sequence

$$Y(n, k) := \binom{n}{k} \frac{sk}{T} T! \sum_{j \geq T} B_{j,T}(a_1, a_2, \dots) D_{z=\beta j+\lambda T}^{ju+vT} \{e^{\alpha z} f_{n-k}(z; a)\} \frac{x^j}{j!} \quad (16)$$

satisfies (4). For $\alpha = 0$ the above sequence can be replaced by

$$Y(n, k) := \frac{n!}{k!} \frac{sk}{T} T! \sum_{j=T}^{h+T} B_{j,T}(a_1, a_2, \dots) \frac{D_{z=\beta j+\lambda T}^{ju+vT} f_{(u+v)T+n-k}(z; a)}{((u+v)T+n-k)!} \frac{x^j}{j!} \quad (17)$$

where $h = \left\lceil \frac{n-k}{u} \right\rceil$ for $u \geq 1$, $h = \infty$ for $u = 0$ and $[x]$ is the largest integer $\leq x$.

More generally, Theorem 3 can be generalized as follows:

Theorem 5 Let $(a_n; n \geq 1)$ be a real sequence, n, r, s, u, v be natural numbers with $r \geq 1$ and $a, \alpha, \beta, \lambda$ be real numbers. Then for $\alpha \neq 0$, the sequence

$$Z(n, s) := \frac{R!}{\gamma^s R} \sum_{j \geq R} B_{j,R}(a_1, a_2, \dots) D_{z=\beta j+\lambda R}^{ju+vR} \{e^{\alpha z} f_n(z; a)\} \frac{x^j}{j!} \quad (18)$$

$$\text{with } \gamma := \alpha^v \varphi(x \alpha^u) \quad \text{and} \quad \varphi(x) := \sum_{i=1}^{\infty} a_i \frac{x^i}{i!}$$

satisfies (7). For $\alpha = 0$ the above sequence can be replaced by

$$Z(n, s) := \frac{n! R!}{\gamma^s R} \sum_{j=R}^{g+R} B_{j,R}(a_1, a_2, \dots) \frac{D_{z=\beta j+\lambda R}^{ju+vR} f_{n+(u+v)R}(z; a)}{(n+(u+v)R)!} \frac{x^j}{j!} \quad (19)$$

$$\text{with } \gamma := \begin{cases} a_1 x (Df_1(0))^{u+v} & \text{if } u \geq 1 \\ (Df_1(0))^v \varphi(x) & \text{if } u = 0 \end{cases} \quad \text{and} \quad \varphi(x) := \sum_{i=1}^{\infty} a_i \frac{x^i}{i!},$$

where $g := \left\lceil \frac{n}{u} \right\rceil$ for $u \geq 1$, and, $g = \infty$ for $u = 0$.

Remark 6 For $a_n = 0$ ($n \geq 2$) in Theorem 4 we obtain Theorem 2.

For $a_n = 0$ ($n \geq 2$) in Theorem 5 we obtain Theorem 3.

For $x = \alpha = 0$ in Theorem 2 we obtain Corollary 5 in [4].

For $x = \alpha = 0$ in Theorem 3 we get Corollary 9 in [4].

From (12) the sequence $Y_1(n, k) := \lim_{\alpha \rightarrow 0} \alpha^{-T} Y(n, k)$ satisfies (4) and gives Proposition 1 in [4], and similarly, from (14) the sequence $Z_1(n, s) := \lim_{\alpha \rightarrow 0} \alpha^{-nr} Z(n, s)$ satisfies (7) and gives Proposition 3 in [4].

By using (8) and (9) we can construct several binomial type polynomials as

$$p_n(t) := t \sum_{k=1}^n Y(n, k) (bn+t)^{k-1}, \quad (p_0(t) := 1),$$

where $(Y(n, k))$ is given by (13) or (12).

3 Applications

We give in this section another versions of Theorems 2, 3 and we present some particular cases of the above results.

3.1 Some applications of Theorem 2

The following corollaries gives a practical version of Theorem 2.

Corollary 7 *Under the hypothesis of Theorem 2 the sequence*

$$Y(n, k) := \binom{n}{k} \frac{sk}{T} \sum_{j=0}^{n-k} \binom{T}{j} D_{z=Tx}^j f_{n-k}(z; a) \alpha^j \quad (20)$$

satisfies (4).

Proof. From (12), we have

$$\begin{aligned} Y(n, k) &= \binom{n}{k} \frac{sk}{T} \sum_{j=0}^{n-k} \binom{T}{j} D_{z=0}^{T-j} \left(e^{z/\alpha} \right) D_{z=0}^j f_{n-k}(Tx + z; a) \\ &= \binom{n}{k} \frac{sk}{T} \sum_{j=0}^{n-k} \binom{T}{j} D_{z=0}^j f_{n-k}(Tx + z; a) \left(\frac{1}{\alpha} \right)^{T-j}. \end{aligned}$$

We conclude noticing that $Y_1(n, k) := \alpha^T Y(n, k)$ satisfies (4). ■

Example 8 *For $f_n(x) = x^n$ the sequence $(Y(n, k))$ in (20) becomes*

$$Y(n, k) := \binom{n}{k} \frac{sk}{T} \sum_{j=0}^{n-k} \binom{T}{j} \frac{(n-k)!}{(n-k-j)!} (Tx + a(n-k))^{n-k-j-1} (Tx + aj) \alpha^j$$

and for $a = 0, \alpha = 1$ the last sequence becomes

$$Y(n, k) := T! \binom{n}{k} \frac{sk}{T} \sum_{j=0}^{n-k} \binom{n-k}{j} \frac{(Tx)^{n-k-j}}{(T-j)!}.$$

The following corollary gives a practical version of Theorem 2 when $\alpha = 0$.

Corollary 9 *Under the hypothesis of Theorem 2 the sequence*

$$\begin{aligned} Y(n, k) &:= \frac{sk}{T} \binom{n}{k} \binom{T+n-k}{n-k} \sum_{j=0}^{-1n-k} \binom{T+j-1}{T-1} B(T+n-k, T+j) \times \\ &\quad (((r+1)c-b)j + b(n-k) + csk)(b(n-k) + csk)^{T+j-1} \end{aligned} \quad (21)$$

satisfies (4), and in particular the sequence

$$Y(n, k) : = \frac{sk}{T} \binom{n}{k} \binom{T+n-k}{n-k}^{-1} \times \sum_{j=0}^{n-k} \binom{T+j-1}{T-1} B(T+n-k, T+j) (T+n-k)^j x^j \quad (22)$$

satisfies (4), where $B_{n,k}(x_1, x_2, x_3, \dots) := B(n, k)$.

Proof. Let $(x_n; n \geq 1)$ be a sequence of real numbers. From (13), it suffices to express $Y(n, k)$ by considering the binomial type sequence

$$f_n(x; a) := x \sum_{k=1}^n B_{n,k}(x_1, x_2, \dots) (an + x)^{k-1} \text{ with } f_0(x) = 1, \quad (23)$$

put after $b := ra + rx + a$, $c := x + a$. To obtain (22), it suffices to choice $b = (r+1)c$, $c = x$ in (21). To obtain (22), it suffices to choice $b = (r+1)c$, $c = x$ in (21). ■

Example 10 By using the well-known identity $B_{n,k}(1, 2, 3, \dots) = \binom{n}{k} k^{n-k}$, if $x_n = n$, the sequence given by (22) becomes

$$Y(n, k) = \frac{n!}{k!} \frac{sk}{(T+n-k)!} \sum_{j=0}^{n-k} \binom{T+n-k}{T+j} (T+j)^{n-k-j} (T+n-k)^j \frac{x^j}{j!},$$

and by using the well-known identity $B_{n,k}(1!, 2!, 3!, \dots) = \binom{n}{k} \frac{(n-1)!}{(k-1)!}$ (the Lah numbers), if $x_n = n!$, the sequence given by (22) becomes

$$Y(n, k) = \frac{n!}{k!} \frac{sk}{T+n-k} \sum_{j=0}^{n-k} \binom{T+n-k}{T+j} (T+n-k)^j \frac{x^j}{j!}.$$

More applications can be constructed by choosing in (21) or (22) x_n as 1, $nf_{n-1}(y; a)$, $D_x f_n(x; a)$.

Example 11 For $r = 0$, $s = 1$, $\alpha = 0$ and $f_n(x) := B_n(x) = \sum_{j=1}^n \binom{n}{j} S(n, j)$ in (13), with $S(n, j)$ are the Stirling numbers of the second kind, we obtain:

$$\begin{aligned} B_{n,k} \left(1, \dots, \frac{x B_{m+1}(am+x) - (x^2 + amx - am) B_m(am+x)}{(am+x)^2}, \dots \right) \\ = kx \sum_{l=0}^{n-k} \binom{l+k}{k} S(n, l+k) (an+kx)^{l-1}. \end{aligned}$$

For $r = 0$, $s = 1$, $\alpha = 0$ and $f_n(x) := [x]_n = \sum_{j=1}^n s(n, j) x^j$ in (13), where $s(n, j)$ are the Stirling numbers of the first kind, we obtain:

$$\begin{aligned} B_{n,k} \left(1, \dots, [am+x-1]_{m-1} \left(m - \sum_{i=1}^{m-1} \frac{am-i}{am+x-i} \right), \dots \right) = \\ kx \sum_{l=0}^{n-k} \binom{l+k}{k} s(n, l+k) (an+kx)^{l-1}. \end{aligned}$$

For $r = 0$, $s = 1$, $\alpha = 0$ and $f_n(x) := [x]^n = \sum_{j=1}^n |s(n, j)| x^j$ in (13), where $|s(n, j)|$ are the absolute Stirling numbers of the first kind, we obtain:

$$\begin{aligned} B_{n,k} \left(1, \dots, [am + x - 1]^{m-1} \left(m - \sum_{i=1}^{m-1} \frac{am + i}{am + i + x} \right), \dots \right) \\ = kx \sum_{l=0}^{n-k} \binom{l+k}{k} |s(n, l+k)| (an + kx)^{l-1}. \end{aligned}$$

3.2 Some applications of Theorem 3

The following corollaries gives a practical version of Theorem 3

Corollary 12 Under the hypothesis of Theorem 3 the sequence

$$Z(n, s) := \frac{1}{R} \sum_{j=0}^n \binom{R}{j} D_{z=Rx}^j f_n(z; a) \alpha^j \quad (24)$$

satisfies (7).

Proof. From (14) we have

$$\begin{aligned} Z(n, s) &= \frac{\alpha^s}{R} D_{z=0}^R \left(e^{z/\alpha} f_n(Rx + z; a) \right) \\ &= \frac{\alpha^s}{R} \sum_{j=0}^n \binom{R}{j} D_{z=0}^j (f_n(Rx + z; a)) \frac{1}{\alpha^{R-j}} \\ &= \frac{1}{R} \sum_{j=0}^n \binom{R}{j} D_{z=0}^j (f_n(Rx + z; a)) \frac{\alpha^j}{\alpha^{nr}}. \end{aligned}$$

We conclude noticing that $Z_1(n, s) := \alpha^{nr} Y(n, k)$ satisfies (4). ■

Example 13 For $f_n(x) = x^n$ and $r \geq 1$ the sequence given by (24) becomes

$$Z(n, s) = \frac{1}{R} \sum_{j=0}^n \binom{R}{j} \frac{n!}{(n-j)!} (Rx + an)^{n-j-1} (Rx + aj) \alpha^j$$

and for $a = 0$, $x = 1$, the last sequence becomes

$$Z(n, s) = \frac{1}{R} \sum_{j=0}^n \binom{R}{j} \frac{n!}{(n-j)!} R^{n-j} \alpha^j.$$

Corollary 14 Under the hypothesis of Theorem 3 the sequence

$$\begin{aligned} Z(n, s) &:= \frac{1}{(Df_1(0))^s} \frac{1}{R} \binom{R+n}{n}^{-1} \sum_{j=0}^n \binom{R+j-1}{R-1} B(R+n, R+j) \\ &\quad \times (cn + bs)^{j-1} \{((r+1)b - c)j + cn + bs\} \end{aligned} \quad (25)$$

satisfies (7), and in particular the sequence

$$Z(n, s) : = \frac{1}{(Df_1(0))^s} \frac{1}{R} \binom{R+n}{n}^{-1} \sum_{j=0}^n \binom{R+j-1}{R-1} \times \quad (26)$$

$$B(R+n, R+j) (R+n)^j x^j$$

satisfies (7), where $B_{n,k}(x_1, x_2, x_3, \dots) := B(n, k)$.

Proof. Let $(x_n; n \geq 1)$ be a sequence of real numbers. From (15), it suffices to express $Z(n, s)$ by considering the binomial type sequence defined in (23), put after $b = a + x$, $c = a + ar + xr$. To obtain (26), it suffices to choice $b = (r+1)c$, $c = x$ in (25). ■

Example 15 If $x_n = n$, the sequence given by (26) becomes

$$Z(n, s) = \frac{1}{(Df_1(0))^s} \sum_{j=0}^n \binom{n}{j} (R+j)^{n-j-1} (R+n)^j x^j$$

and if $x_n = n!$, the sequence given by (26) becomes

$$Z(n, s) = \frac{(R+n-1)!}{(Df_1(0))^s} \sum_{j=0}^n \binom{n}{j} \frac{(R+n)^j}{(R+j)!} x^j.$$

3.3 Some applications of Theorem 4

Some particular cases of Theorem 4 are given by the following corollaries:

Corollary 16 Under the hypothesis of Theorem 4 and $v \geq u$ the sequences

$$Y_1(n, k) : = \binom{n}{k} \frac{sk}{T} \sum_{j=0}^T \binom{T}{j} D_{z=\beta j + \lambda T}^{ju+VT} \{e^{\alpha z} f_{n-k}(z; a)\} x^j y^{T-j}, \quad (27)$$

$$Y_2(n, k) : = \frac{n!}{k!} \frac{sk}{T} \sum_{j=0}^T \binom{T}{j} \frac{D_{z=\beta j + \lambda T}^{ju+VT} f_{n-k}(z; a)}{(vT + n - k)!} x^j y^{T-j} \quad (28)$$

satisfy (4).

Proof. Let $a_1 = p$, $a_2 = 2q$ and $a_m = 0$ for $m \geq 3$. To obtain (27) it suffices to replace the identity $B_{j,T}(p, 2q, 0, 0, \dots) = \binom{T}{j-T} p^{2T-j} q^{j-T}$ in (17) and in (16). ■

Some particular cases of (29) are given by:

For $r = u = 0$, $v = 1$ or $u = v = r = 0$ in (27), the sequences

$$Y_1(n, k) : = \binom{n}{k} \sum_{j=0}^{sk} \binom{sk}{j} D_{z=\beta j + \lambda k}^{sk} \{e^{\alpha z} f_{n-k}(z; a)\} x^j y^{sk-j}, \quad (29)$$

$$Y_2(n, k) : = \frac{n!}{k!} \sum_{j=0}^{sk} \binom{sk}{j} \frac{D_{z=\beta j + \lambda k}^{sk} f_{n+(s-1)k}(z; a)}{(n + (s-1)k)!} x^j y^{sk-j}, \quad (30)$$

$$Y_3(n, k) : = \binom{n}{k} \sum_{j=0}^{sk} \binom{sk}{j} f_{n-k}(\beta j + \lambda k; a) x^j y^{sk-j} \quad (31)$$

satisfy (4), and if $s = 1$ the last sequences give

$$\begin{aligned} & B_{n,k}(\alpha y, \dots, m(\alpha y f_{m-1}(\lambda; a) + x D_{z=\beta} f_{m-1}(z; a)), \dots) \\ &= \binom{n}{k} \sum_{j=0}^k \binom{k}{j} D_{z=0}^k \{e^{\alpha z} f_{n-k}((\beta - \lambda)j + \lambda k + z; a)\} x^j y^{k-j}, \end{aligned} \quad (32)$$

$$\begin{aligned} & B_{n,k}((x+y) D f_1(0), \dots, y D_{z=\lambda} f_n(z; a) + x D_{z=\beta} f_n(z; a), \dots) \\ &= \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} D_{z=0}^k f_n((\beta - \lambda)j + \lambda k + z; a) x^j y^{k-j} \end{aligned} \quad (33)$$

$$\begin{aligned} & B_{n,k}(y+x, \dots, m(y f_{m-1}(\lambda; a) + x f_{m-1}(\beta; a)), \dots) \\ &= \binom{n}{k} \sum_{j=0}^k \binom{k}{j} f_{n-k}((\beta - \lambda)j + \lambda k; a) x^j y^{k-j} \end{aligned} \quad (34)$$

Example 17 For $f_n(x) = x^n$, $a = 0$ and $u = 0$ or 1 in (27) the sequences

$$\begin{aligned} Y_1(n, k) &= \frac{sk}{T} \binom{n}{k} \sum_{j=0}^T \binom{T}{j} (\alpha T + \beta j)^{n-k} x^{T-j} y^j \text{ and} \\ Y_2(n, k) &= \frac{n!}{k!} \frac{sk}{T} \sum_{j=0}^{n-k} \binom{T}{j} \frac{(\alpha T + \beta j)^{n-k-j}}{(n-k-j)!} x^{T-j} y^j \end{aligned}$$

satisfy (4), and for $f_n(x) = x^n$, $a = 0$, $u = 0$ or 1 , $a_n = n$ in Theorem 4 the sequences

$$\begin{aligned} Y_3(n, k) &= \frac{sk}{T} \binom{n}{k} \sum_{j=0}^{\infty} (\alpha T + \beta j)^{n-k} \frac{(Tx)^j}{j!} \text{ and} \\ Y_4(n, k) &= \frac{sk}{T} \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (\alpha T + \beta j)^{n-k-j} (Tx)^j \end{aligned}$$

satisfy (4).

An interesting relation between Bell polynomials and Appell polynomials can be viewed as a special case of Theorem 4 and it is given by:

Corollary 18 For the sequence of polynomials $(A_n(x, y, z))$ defined by

$$A_n(x, y, z) := \sum_{j=0}^n \binom{n}{j} a_j ((x+y)j + x + y + z)(xj + yn + x + y + z)^{n-1-j}$$

we have

$$\begin{aligned} & B_{n,k}(A_0(x, y, z), \dots, mA_{m-1}(x, y, z), \dots) \\ &= \sum_{j=k}^n \binom{n}{j} B_{j,k}(a_0, 2a_1, 3a_2, \dots) (jx + ny + kz)^{n-j-1} ((x+y)j + kz), \end{aligned} \quad (35)$$

and in particular when $x = y = 0$ we get

$$B_{n,k}(A_0(z), \dots, mA_{m-1}(z), \dots) = \sum_{j=k}^n \binom{n}{j} B_{j,k}(a_0, \dots, ma_{m-1}, \dots) (kz)^{n-j}, \quad (36)$$

where $A_n(z) := \sum_{j=0}^n \binom{n}{j} a_j z^{n-j}$ is an Appell polynomial.

Proof. It suffices to use (17) with $x = 1$, $r = v = 0$, $s = u = 1$, $f_n(x; a) := a(an + x)^{n-1}$ and replace a_n by na_{n-1} . ■

Example 19 For any sequence $(\varphi_n; n \geq 1)$ for real numbers, let I_n be the identity matrix of order n and (A_n) be the sequence of matrices defined by: $A_0 := 1$, $A_n := (a_{ij})$ for $1 \leq i, j \leq n$ with $a_{ij} = \varphi_{j-i+1}$ if $j \geq i$, $a_{i,i-1} = i - 1$ and $a_{ij} = 0$ otherwise. Then from [7, p. 110] and from (36) we get:

$$B_{n,k}(1, \dots, m \det(A_{m-1} + xI_{m-1}), \dots) = \sum_{j=k}^n \binom{n}{j} B_{j,k}(1, \dots, m \det A_{m-1}, \dots) (kx)^{n-j}.$$

3.4 Some applications of Theorem 5

A particular case of Theorem 5 is given by the following corollary:

Corollary 20 Under the hypothesis of Theorem 5 and $v \geq u$ the sequences

$$Z_1(n, s) : = \frac{1}{\gamma_1^s R} \sum_{j=0}^R \binom{R}{j} D_{z=\beta j + \lambda R}^{ju+vR} \{e^{\alpha z} f_n(z; a)\} x^j y^{R-j} \text{ and} \quad (37)$$

$$Z_2(n, s) : = \frac{n!}{\gamma_2^s R} \sum_{j=0}^R \binom{R}{j} \frac{D_{z=\beta j + \lambda R}^{ju+vR} f_{n+(u+v)R}(z; a)}{(n + (u+v)R)!} x^j y^{R-j} \quad (38)$$

satisfy (7), where $\gamma_1 := \alpha^v (x\alpha^u + y)$ and $\gamma_2 := \begin{cases} (x+y)(Df_1(0))^v & \text{if } u \geq 1 \\ y(Df_1(0))^v & \text{if } u = 0 \end{cases}$.

Proof. It suffices to put in Theorem 5 $x = 1$; $a_1 = p$, $a_2 = 2q$ and $a_m = 0$ for $m \geq 3$ and use the identity $B_{j,R}(p, 2q, 0, 0, \dots) = \binom{R}{j-R} p^{2R-j} q^{j-R}$. ■

Example 21 For $f_n(x) = x^n$, $a = 0$ and $u = 0$ or 1 the sequence given by (37) becomes

$$Z_1(n, s) = (x+y)^{-s} \frac{1}{R} \sum_{j=0}^R \binom{R}{j} (\beta j + \alpha R)^n x^j y^{R-j} \text{ and}$$

$$Z_2(n, s) = \frac{n!}{R} \sum_{j=0}^n \frac{1}{(n-j)!} \binom{R}{j} (\beta j + \alpha R)^{n-j} x^j$$

satisfy (7), and for $f_n(x) = x^n$, $a = 0$, $x = 1$, $u = 0$ or 1 , $a_n = nt^{n-1}$ the sequence given by (19) becomes

$$Z_3(n, s) = \frac{1}{R} \sum_{j=0}^n \binom{n}{j} (\beta j + \alpha R)^{n-j} R^j t^j \text{ and}$$

$$Z_4(n, s) = \frac{\exp(-sx)}{R!} \sum_{j=0}^{\infty} (\beta j + \alpha R)^n \frac{(Rx)^j}{j!}.$$

4 Proof of the main results

Proof of Lemma 1. Let $F(t; a)^x := 1 + \sum_{n=1}^{\infty} f_n(x; a) \frac{t^n}{n!}$. Now, because $f_n(x; a)$ is a polynomial of degree n , then the proof follows from the following expansions:

$$\sum_{n=0}^{\infty} D_{z=0}^k (e^{\alpha z} f_n(kx + z; a)) \frac{t^{n+k}}{n!} = \sum_{n=k}^{\infty} \frac{n!}{(n-k)!} D_{z=0}^k (e^{\alpha z} f_{n-k}(kx + z; a)) \frac{t^n}{n!}$$

and

$$\begin{aligned} \sum_{n=0}^{\infty} D_{z=0}^k (e^{\alpha z} f_n(kx + z; a)) \frac{t^{n+k}}{n!} &= D_{z=0}^k \left(e^{\alpha z} \sum_{n=0}^{\infty} f_n(kx + z; a) \frac{t^{n+k}}{n!} \right) \\ &= D_{z=0}^k \left(e^{\alpha z} F(t; a)^{kx+z} t^k \right) \\ &= t^k F(t; a)^{kx} D_{z=0}^k \left(e^{(\alpha + \ln F(t; a))z} \right) \\ &= t^k F(t; a)^{kx} (\alpha + \ln F(t; a))^k \\ &= t^k e^{-\alpha kx} ((e^{\alpha} F(t; a))^x \ln(e^{\alpha} F(t; a)))^k \\ &= t^k e^{-\alpha kx} (D_x(e^{\alpha} F(t; a))^x)^k \\ &= t^k \left(\sum_{m=0}^{\infty} e^{-\alpha x} D_x(e^{\alpha x} f_m(x; a)) \frac{t^m}{m!} \right)^k \\ &= \left(\sum_{m=1}^{\infty} m e^{-\alpha x} D_x(e^{\alpha x} f_{m-1}(x; a)) \frac{t^m}{m!} \right)^k \\ &= k! \sum_{n=k}^{\infty} B_{n,k}(\alpha, \dots, m e^{-\alpha x} D_x(e^{\alpha x} f_{m-1}(x; a)), \dots) \frac{t^n}{n!}. \end{aligned}$$

Then by identification we get

$$B_{n,k}(\alpha, 2e^{-\alpha x} D_x(e^{\alpha x} f_1(x; a)), \dots, m D_x(e^{\alpha x} f_{m-1}(x; a))) = \binom{n}{k} D_{z=0}^k (e^{\alpha z} f_{n-k}(kx + z; a)).$$

To obtain (10), it suffices to remark that for $m \geq 1$ we have

$$e^{-\alpha x} D_x(e^{\alpha x} f_m(x; a)) = D_{z=0}(e^{\alpha z} f_m(x + z; a)).$$

The identity (10) can be written when $\alpha = 0$ as

$$B_{n,k}(0, \dots, m D_{z=0} f_{m-1}(kx + z; a), \dots) = \binom{n}{k} D_{z=0}^k f_{n-k}(kx + z; a)$$

which is equivalent to

$$B_{n,k}(D_x f_1(x; a), \dots, D_x f_m(x; a), \dots) = \frac{1}{k!} D_{z=0}^k f_n(kx + z; a).$$

■

Proof of Theorem 2. When we replace x_n by αx_n in (2) and we use the well-known identities

$$B_{n,k}(\alpha x_1, \alpha x_2, \alpha x_3, \dots) = \alpha^k B_{n,k}(x_1, x_2, x_3, \dots) \quad (39)$$

$$\text{and } B_{n,k}(\alpha x_1, \alpha^2 x_2, \alpha^3 x_3, \dots) = \alpha^n B_{n,k}(x_1, x_2, x_3, \dots), \quad (40)$$

it results that the identity (2) remains true for $x_1 \neq 1$. Then for $r + s \geq 1$ and for the choice $x_n = nD_{z=0}(e^{\alpha z} f_{n-1}(x+z; a))$ in (2), the identity (10) proves that the sequence $(Y(n, k))$ given by (3) becomes

$$\begin{aligned} Y(n, k) &= \binom{n}{k} \frac{sk}{T} \binom{T+n-k}{T}^{-1} B_{T+n-k, T}(\alpha, \dots, mD_{z=0}(e^{\alpha z} f_{m-1}(x+z; a)), \dots) \\ &= \binom{n}{k} \frac{sk}{T} D_{z=0}^T \{e^{\alpha z} f_{n-k}(Tx+z; a)\}. \end{aligned}$$

For the particular case $\alpha = 0$, if we take $x_n = D_{z=0}(f_n(x+y; a))$ in (2) and we use the identity (11), the sequence $(Y(n, k))$ given by (3) becomes

$$\begin{aligned} Y(n, k) &= \binom{n}{k} \frac{sk}{T} \binom{T+n-k}{T}^{-1} B_{T+n-k, T}(D_{z=0}(e^{\alpha z} f_1(x+z; a)), \dots) \\ &= \frac{n!}{k!} \frac{sk}{T} \frac{1}{(T+n-k)!} D_{z=0}^T f_{T+n-k}(Tx+z; a). \end{aligned}$$

Note that for the case $r = s = 0$, the sequence $(Y(n, k))$ given by (12) is not defined. We put in this case $Y(n, k) = \binom{n}{k} f_{n-k}(x; a)$. Proposition 1 in [4] proves that this sequence satisfies (4). ■

Proof of Theorem 3. Case $\alpha \neq 0$: Let $x_n = \frac{n}{\alpha} D_{z=0} \{e^{\alpha z} f_{n-1}(x+z; a)\}$. We have $x_1 = 1$ and then by using the identity (10), the sequence $(Z(n, s))$ given by (6) becomes

$$\begin{aligned} Z(n, s) &= \frac{1}{\alpha^R} \frac{1}{R} \binom{R+n}{R}^{-1} B_{R+n, R}(\alpha, \dots, mD_{z=0}(e^{\alpha z} f_{m-1}(x+z; a)), \dots) \\ &= \frac{1}{\alpha^R} \frac{1}{R} D_{z=0}^R (e^{\alpha z} f_n(Rx+z; a)). \end{aligned}$$

Case $\alpha = 0$: Let $x_n = D_x f_n(x; a) / D_x f_1(x; a)$. Now because $D_x f_1(x) = x D f_1(0)$ we get $D_x f_1(x; a) = D_x \left(\frac{x}{x+a} f_1(x+a) \right) = D f_1(0) \neq 0$. We have $x_1 = 1$ and then by using the identity (11), the sequence $(Z(n, s))$ given by (6) becomes

$$\begin{aligned} Z(n, s) &= \frac{1}{R} \binom{R+n}{R}^{-1} \frac{B_{R+n, R}(D_{z=0}(f_1(x+z; a)), \dots, D_{z=0}(f_m(x+z; a)), \dots)}{(D_x f_1(x; a))^R} \\ &= \frac{1}{(D f_1(0))^R} \frac{1}{R} \frac{n!}{(R+n)!} D_{z=0}^R f_{R+n}(kx+z; a). \end{aligned}$$

Note that if $Z(n, s)$ satisfies (7) then $\lambda^n Z(n, s)$ satisfies (7). ■

Proof of Theorem 4. Let $(a_n; n \geq 1)$ be a real sequences; u, v be a natural numbers and we put $F(t)^x := 1 + \sum_{n=1}^{\infty} f_n(x) \frac{t^n}{n!}$ with $f_0(x) := 1$.

For $F(n, k) := \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) D_{z=0}^{ju+vk} \{e^{\alpha z} f_n(\beta j + \lambda k + z; a)\} \frac{x^j}{j!}$ we have

$$\begin{aligned}
\sum_{n \geq 0} F(n, k) \frac{t^{n+k}}{n!} &= t^k \sum_{n \geq 0} \frac{t^n}{n!} \left(\sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) D_{z=0}^{ju+vk} \{e^{\alpha z} f_n(\beta j + \lambda k + z; a)\} \frac{x^j}{j!} \right) \\
&= t^k \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) \frac{x^j}{j!} \left(\sum_{n \geq 0} D_{z=0}^{ju+vk} \{e^{\alpha z} f_n(\beta j + \lambda k + z; a)\} \frac{t^n}{n!} \right) \\
&= t^k \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) \frac{x^j}{j!} D_{z=0}^{ju+vk} \left\{ e^{\alpha z} \sum_{n \geq 0} f_n(\beta j + \lambda k + z; a) \frac{t^n}{n!} \right\} \\
&= t^k \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) \frac{x^j}{j!} D_{z=0}^{ju+vk} \left\{ e^{\alpha z} (F(t; a))^{\beta j + \lambda k + z} \right\} \\
&= t^k \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) \frac{x^j}{j!} (F(t; a))^{\beta j + \lambda k} D_{z=0}^{ju+vk} \{e^{\alpha z} F(t; a)^z\} \\
&= t^k \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) \frac{x^j}{j!} (F(t; a))^{\beta j + \lambda k} (\alpha + \ln F(t; a))^{ju+vk} \\
&= t^k \sum_{j \geq k} B_{j,k}(u_1, u_2, \dots) \frac{x^j}{j!} \\
&= \frac{t^k}{k!} \left(\sum_{m \geq 1} u_m \frac{x^m}{m!} \right)^k \\
&= \frac{t^k}{k!} \left(\sum_{m \geq 1} a_m F(t; a)^{\beta m + \lambda} (D_{z=0} (e^{\alpha z} (F(t; a))^z))^{mu+v} \frac{x^m}{m!} \right)^k \\
&= \frac{t^k}{k!} \left(\sum_{m \geq 1} a_m D_{z=0}^{mu+v} (e^{\alpha z} F(t; a)^{\beta m + \lambda + z}) \frac{x^m}{m!} \right)^k \\
&= \frac{t^k}{k!} \left(\sum_{m \geq 1} a_m \sum_{j \geq 0} D_{z=0}^{mu+v} \left(e^{\alpha z} f_j(\beta m + \lambda + z; a) \frac{t^j}{j!} \right) \frac{x^m}{m!} \right)^k \\
&= \frac{t^k}{k!} \left(\sum_{j \geq 0} \frac{t^j}{j!} \sum_{m \geq 1} a_m D_{z=0}^{mu+v} (e^{\alpha z} f_j(\beta m + \lambda + z; a)) \frac{x^m}{m!} \right)^k \\
&= \frac{t^k}{k!} \left(\sum_{j \geq 0} v_j \frac{t^j}{j!} \right)^k \\
&= \frac{1}{k!} \left(\sum_{j \geq 1} j v_{j-1} \frac{t^j}{j!} \right)^k \\
&= \sum_{n \geq k} B_{n,k}(v_0, \dots, j v_{j-1}, \dots) \frac{t^n}{n!},
\end{aligned}$$

where $u_m = a_m (F(t; a))^{\beta m + \lambda} (\alpha + \ln F(t; a))^{mu+v} = a_m F(t; a)^{\beta m + \lambda} (D_{z=0} (e^{\alpha z} (F(t; a))^z))^{\mu u+v}$ ($m \geq 1$) and $v_m = \sum_{m \geq 1} a_m D_{z=0}^{\mu u+v} (e^{\alpha z} f_j(\beta m + \lambda + z; a)) \frac{x^m}{m!} = F(m, 1)$ ($m \geq 0$). Then, we obtain

$$B_{n,k}(F(0, 1), \dots, jF(j-1, 1), \dots) = k! \binom{n}{k} F(n-k, k)$$

and this means that the sequence

$$X(n, k) := k! \binom{n}{k} \sum_{j \geq k} B_{j,k}(a_1, a_2, \dots) D_{z=0}^{ju+vk} \{e^{\alpha z} f_{n-k}(\beta j + \lambda k + z; a)\} \frac{x^j}{j!} \quad (41)$$

satisfies (4). To obtain (16), it suffices to take $x_n = X(n, 1)$ in (2), where $(X(n, k))$ is the sequence given by (41). Indeed, the sequence given by (3) becomes

$$\begin{aligned} Y(n, k) &= \binom{n}{k} \frac{sk}{T} \binom{T+n-k}{T}^{-1} B_{T+n-k, T}(X(1, 1), X(2, 1), \dots) \\ &= \binom{n}{k} \frac{sk}{T} \binom{T+n-k}{T}^{-1} X(T+n-k, T) \\ &= \binom{n}{k} \frac{sk}{T} T! \sum_{j \geq T} B_{j,T}(a_1, a_2, \dots) D_{z=0}^{ju+vT} \{e^{\alpha z} f_{n-k}(\beta j + \lambda T + z; a)\} \frac{x^j}{j!}. \end{aligned}$$

For the particular case $\alpha = 0$ and $u \geq 1$ we remark that $X(n, 1) = 0$ for $u + v > n - 1$ and then the identity (41) becomes

$$B_{n,k} \left(\overbrace{0, \dots, 0}^{u+v}, X(u+v+1, 1), X(u+v+2, 1), \dots \right) = X(n, k) \quad (42)$$

or, equivalently, by using the well-known identity

$$B_{n,k}(0, \dots, 0, a_{r+1}, a_{r+2}, \dots) = \frac{n!}{(n-rk)!} B_{n-rk,k} \left(\frac{a_{1+r}}{(1+r)!}, \dots, \frac{i! a_{i+r}}{(i+r)!}, \dots \right).$$

Setting $X^*(n, k) := n! X(n + (u+v)k, k) / (n + (u+v)k)!$, the identity (42) becomes

$$B_{n,k}(X^*(1, 1), X^*(2, 1), \dots) = X^*(n, k)$$

To obtain (17) it suffices to take $x_n = X^*(n, 1)$ in (2). Indeed

$$\begin{aligned} Y(n, k) &= \binom{n}{k} \frac{sk}{T} \binom{T+n-k}{T}^{-1} B_{T+n-k, T}(X^*(1, 1), X^*(2, 1), \dots) \\ &= \binom{n}{k} \frac{sk}{T} \binom{T+n-k}{T}^{-1} X^*(T+n-k, T) \\ &= \frac{n!}{k!} \frac{sk}{T} T! \sum_{j \geq T} B_{j,T}(a_1, a_2, \dots) D_{z=0}^{ju+vT} \left\{ e^{\alpha z} \frac{f_{(u+v)T+n-k}(\beta j + \lambda T + z; a)}{((u+v)T+n-k)!} \right\} \frac{x^j}{j!}. \end{aligned}$$

■

Proof of Theorem 5. For $\alpha \neq 0$ it suffices to put in (5) $x_n = X(n, 1)/X(1, 1)$, where $(X(n, k))$ is the sequence given by (41). We have

$$x_1 = X(1, 1) = \sum_{j \geq 1} a_j \alpha^{ju+v} \frac{x^j}{j!} = \alpha^v \varphi(x \alpha^u) \quad \text{with } \varphi(x) := \sum_{j \geq 1} a_j \frac{x^j}{j!}.$$

The sequence $(Z(n, s))$ given by (6) becomes

$$\begin{aligned} Z(n, s) &= \frac{1}{(X(1, 1))^R} \frac{1}{R} \binom{R+n}{R}^{-1} X(R+n, R) \\ &= \frac{1}{(\alpha^v \varphi(x \alpha^u))^R} \frac{R!}{R} \sum_{j \geq R} B_{j,R}(a_1, a_2, \dots) D_{z=0}^{ju+vR} \{e^{\alpha z} f_n(\beta j + \lambda R + z; a)\} \frac{x^j}{j!}. \end{aligned}$$

For $\alpha = 0$ it suffices to put in (5) $x_n = X^*(n, 1)/X^*(1, 1)$, where $(X^*(n, k))$ is the sequence given by (??). We have

$$X^*(1, 1) = \frac{X(u+v+1, 1)}{(u+v+1)!} = \begin{cases} a_1 x (Df_1(0))^{u+v} & \text{if } u \geq 1 \\ (Df_1(0))^v \varphi(x) & \text{if } u = 0 \end{cases}$$

The sequence $(Z(n, s))$ given by (6) becomes

$$\begin{aligned} Z(n, s) &= \frac{1}{(X^*(1, 1))^R} \frac{1}{R} \binom{R+n}{R}^{-1} X^*(R+n, R) \\ &= \frac{n!}{(X^*(1, 1))^R} \frac{R!}{R} \sum_{j \geq R} B_{j,R}(a_1, a_2, \dots) D_{z=0}^{ju+vR} \left\{ \frac{f_{n+(u+v)R}(\beta j + \lambda R + z; a)}{(n+(u+v)R)!} \right\} \frac{x^j}{j!}. \end{aligned}$$

■

Acknowledgement 22 *The author thanks the Professor Hacene Belbachir for his careful reading and valuable suggestions.*

References

- [1] M. Abbas and S. Bouroubi, On new identities for Bell's polynomials, Discrete Mathematics 293 (2005) 5-10.
- [2] E.T. Bell, Exponential polynomials, Annals of Mathematics 35, 1934, 258-277.
- [3] L. Comtet, Advanced Combinatorics. D. Reidel Publishing Company, Dordrecht-Holland / Boston-U.S.A, 1974, 133-175.
- [4] M. Mihoubi, Bell polynomials and binomial type sequences. Discrete Mathematics, 308 (2008) 2450 – 2459.
- [5] J. Riordan, Combinatorial Identities. Huntington, New York, 1979.
- [6] S. Roman, The Umbral Calculus. Academic Press, INC, 1984.
- [7] R. Vein and P. Dale, Determinants and their applications in mathematical physics. Springer-Verlag New York, INC, 1999.